

## Native Herbaceous Plants for Sun to Part Sun

Manada Native Plant Database					
Scientific Name	Common name	Bloom Time/Fruiting Time	Bloom Color	Soil Moisture	Height
Agastache nepetoides (H)	Giant Yellow Hyssop	Late Summer	Light Yellow	Moist to Dry, well-drained	3'-8'/2'-3'
Agastache scrophulariifolia (H)	Purple-Giant Hyssop	Late Summer	Light Pink or Lavendar	Moist to Dry	3'-6'/2'-3'
Allium cernuum (H)	Nodding Wild Onion	Summer	Pink	Moist to Dry	1'-1.5'/1'
Amsonia tabermontana (H)	Eastern Blue Star	Late Spring	Light Blue	Moist to Dry	3'-3.5'/3'
Anaphalis margaritacea (H) **	Pearly-Everlasting	Summer	White	Dry	1'-3'/1'-2'
Anemone canadensis (H)	Canada Anemome	Late Spring	White	Moist to Dry	1'-2'/3'-6'
Anemone virginiana (H)	Tall Anemome, Tall Thimbleweed,	Summer	White	Moist to Dry	12"-30"/12"-16"
Antennaria plantaginifolia (H)	Plantain-leaved Pussytoes	Spring	White, Fuzzy	Moist to Dry	3"-6"/24"
Aquilegia canadensis (H)	Wild Columbine	Spring	Red & Yellow	Moist to Dry	1'-2'/1'-2'
Asclepias incarnata (H)	Swamp Milkweed	Summer	Pink	Wet to Moist to Average	2'-4'/2'
Asclepias pururascens (H)	Purple Milkweed	Summer	Rose Purple	Moist to Dry	2' - 4'/2'
Asclepias syriaca (H)	Common Milkweed	Summer - June, July	Pink and light purple	Moist to Dry	2'-4"/3'-5'
Asclepias tuberosa (H) **	Butterfly Milkweed	Summer - June, July	Orange	Dry to Average well drained	1'-3'/2'
Asclepias verticillata (H) **	Whorled Milkweed, Horsetail Milkweed	Summer - June, July	White	Moist to Dry	1'-2'/1.5'
Aster ericoides (Symphyotrichum ericoides var. ericoides) (H)	Heath Aster	Late Summer - Fall	White	Moist to Dry	1'-3'/2'
Aster laevis (Symphyotrichum) (H)	Smooth Aster	Fall - September, October	Violet	Moist to Dry	2' - 4'/12"-16"
Aster lateriflorus (H)	Calico Aster	Late Summer-Fall	White with reddish or yellow centers	Average to Dry	1'-5'
Aster novae-angliae (Symphyotrichum) (H)	New England Aster	Fall	Puple	Moist to Average	2'-6'/2'-3'
Aster novae-belgi (Symphyotrichum novi-belgii var. novi-belgii) (H)	New York Aster	Fall	Violet, Rose, White	Wet, Moist to Average	1'-3'/2'
Aster oblongifolius	Aromatic Aster	Fall -Late-	Purple, blueish,	Average to Dry	2'-3'/3'

## Native Herbaceous Plants for Sun to Part Sun

Manada Native Plant Database					
Scientific Name	Common name	Bloom Time/Fruiting Time	Bloom Color	Soil Moisture	Height
(Symphyotrichum oblongifolium) (H) **		October, November	pink		
Aster puniceus (Symphyotrichum puniceum var. puniceum) (H) **	Purple-stemmed Aster	Late Summer - Fall	Light purple	Wet to Moist	2'-6'/2'
Aster spectabilis (Eurybia spectabilis) (H) **	Showy Aster	Late Summer - Fall	Violet, Lavendar	Moist to Dry	1'-3'/2'
Baptisia australis (H)	Blue Wild Indigo, Blue False Indigo	Spring into Summer	Indigo Blue	Moist to Dry	3' - 4'/3'
Baptisia tinctoria (H)	Wild Indigo, Wild Yellow Indigo	Late Spring - Summer	Yellow	Moist to Dry	2'-3'/2'-3'
Boltonia asteroides (H) **	False Aster, Thousand Flowered Aster	Late Summer - Early fall	White	Moist to Average	3'-4'/3'
Caltha palustris (H)	Marsh Marigold	Early Spring	Yellow	Wet to Moist	1'-2'/1'
Campanula rotundiflora (H)	Harebell	Summer-Fall	Blue-Violet	Moist to Dry	8"-14"
Chelone glabra (H)	White Turtlehead	Late Summer - Fall	White	Wet, Moist to Average	3'/2'
Chrysopsis mariana (H)	Golden Aster, Maryland Golden Aster	Late summer to early fall	Yellow	Well drained	1'-2.5'/12"
Conoclinium coelestinum (Eupatorium coelestinum) (H)	Mist Flower	Fall	Blue-Violet	Moist to Average	2' - 3'/3'
Coreopsis rosea (H)	Pink Tickseed	Summer	Pink with Yellow Eye	Wet to Moist	1'/1'
Coreopsis tripteris (H) **	Tall Coreopsis	Mid to Late Summer	Yellow	Moist to Average	3'-6'/3/
Dicentra exima (H)	Wild Bleeding Heart	Spring through Fall	Pink	Moist	1'-1.5'/1.5'-2
Dodecatheon meadia (H)	Shooting Star	Spring	Deep Pink	Moist	8"-14"/8"-12"
Eupatoriadelphus dubius (Eupatorium dubium) (H)	Three-nerved Joe-pye weed	Late Summer	Pink	Wet, Moist to Average	3' - 4'
Eupatoriadelphus fistulosus (Eupatorium fistulosum) (H) **	Hollow Joe- Pye Weed, Hollow-stemmed Joe-Pye Weed, Trumpetweed	Late Summer	Pink	Moist to Average	5'-8'/3'-4'

## Native Herbaceous Plants for Sun to Part Sun

Manada Native Plant Database					
Scientific Name	Common name	Bloom Time/Fruiting Time	Bloom Color	Soil Moisture	Height
Eupatoriadelphus maculatus (Eupatorium maculatum) (H) **	Spotted Joe-pye weed	Late Summer	Mauve Pink	Moist to Average	5'-8'/3'-5'
Eupatorium hyssopifolium (H) **	Hyssop-Leaved Eupatorium	Late Summer	White	Dry, well-drained	2'-3'/1'-2'
Eupatorium perfoliatum (H)	Boneset	Late Summer into Fall	White	Wet to Moist	2'-4'/2'
Eupatorium purpurea (H)	Sweet-Scented Joe-Pye weed	Late Summer	Mauve Pink	Moist	5'-7'/2'-4'
Eupatorium rotundifolium (H) **	Round-leaf Thoroughwort, Round-leaved Boneset	Summer into Fall	White	Moist to Dry	2'-5'/2'
Eupatorium sessilifolium (H)	Upland Boneset	Late Summer	White	Moist to Dry	2'-5'/2'
Euthamia graminifolia (Solidago) (H) **	Flat-Top Goldentop	Summer	Yellow	Moist to Dry	2.5'-3'/2'
Filipendula Rubra (H)	Queen of the Prairie	Summer	Pink	Wet to Moist	3'-7'/3'-4'
Gentiana andrewsii var. andrewsii (H) ***	Bottle Gentian	Late Summer-Fall	Deep Blue	Moist	1'-2'/1'
Gentiana clausa (H) ***	Meadow Closed Gentian, Meadow Bottle Gentian	Fall	Pale to Deep Blue/Violet	Wet to Moist	14"-18"/24"-30"
Geranium maculatum (H)	Wild Geranium, Cranesbill	Spring	Pinkish	Moist in spring	14"-24"/12"
Gillenia trifoliata (Porteranthus stipulatus) (H)	Bowman's Root	Late Spring into Summer	White	Moist to Dry	2'-3'/2'
Helenium autumnale (H)	Sneezeweed, Helens flowers	Late Summer	Yellow, Rust	Wet to Moist	3' - 5'/ 3'
Helenium flexuosum (H)	Purplehead Sneezeweed	Summer	Yellow	Moist	1'-3'/1'-1.5'
Helianthus angustifolius	Swamp Sunflower, Narrow-Leaf Sunflower	Fall	Yellow	Wet to Moist	5'-9'/3'-4')
Helianthus decapetalus (H)	Thin-leaved Sunflower	Late Summer	Yellow	Moist to Average	2'-5'/4'-5'
Helianthus divaricatus (H)	Woodland Sunflower	Summer	Yellow	Moist to Moderately Dry	2'-4'/3'-6'
Helianthus microcephalus (H)	Small-headed Sunflower	Late Summer	Yellow	Moist	3'-6'/3'-4'

## Native Herbaceous Plants for Sun to Part Sun

Manada Native Plant Database					
Scientific Name	Common name	Bloom Time/Fruiting Time	Bloom Color	Soil Moisture	Height
<i>Heliopsis helianthoides</i> (H)	Oxeye Sunflower	Summer	Yellow	Moist to Dry	4'-6'/3'-5'
<i>Hibiscus moscheutos</i> (H) (E) **	Common Rose Mallow, Swamp Rose Mallow	Late Summer	White to Pink	Wet to Moist	4'-7'/3'-4'
<i>Houstonia caerulea</i>	Bluelets, Quaker Ladies	Spring	Light Blue	Wet to Moist	1"-4"/3"
<i>Hypoxis hirsuta</i>	Common Star Grass, Yellow Star Grass	Late Spring into Summer	Bright Yellow	Moist to Dry	3"-6"/6"-8"
<i>Impatiens capensis</i> (H)***	Jewelweed, Touch-Me-Not	Late Summer - Fall	Bright Orange	Wet to Moist	2'-5'/2'
<i>Impatiens pallida</i> (H)***	Pale Jewelweed, Pale Touch-Me-Not	Late Summer - Fall	Pale Yellow	Wet to Moist	3'-6'/2'
<i>Iris versicolor</i>	Northern Blue Flag	Late Spring	Blue-Violet, Purple	Wet to Moist	3'-4'/2'-3'
<i>Liatris scariosa</i> ( <i>Liatris borealis</i> ) (H)	Northern Blazingstar	Late Summer	Pink-purple	Moist to Dry	2'-4'/1'-1.5'
<i>Liatris spicata</i> (H) **	Blazing Star, Gayfeather	Summer	Pink-purple	Moist to Dry	2' - 4'/18"
<i>Liatris squarrosa</i> (H) **	Scaly Blazingstar	Mid to Late Summer	Purple	Moist to Dry	24"-30"/12"-15"
<i>Lilium canadense</i> (H)	Canada Lily	Summer	Yellow to Yellow Orange	Moist to Wet, acidic	3'-8'/18"-24"
<i>Lilium superbum</i> (H)	Turk's Cap, Turk's Cap Lily	Summer	Orange with Black Freckles	Moist to Wet	3'-8'/18"-24"
<i>Lobelia cardinalis</i> (H)	Cardinal Flower	Summer	Scarlet	Wet to Moist	2' - 4'/1'
<i>Lobelia siphilitica</i> (H)	Great Blue Lobelia	Summer	Blue	Wet to Dry	2' - 3'/1'-1.5'
<i>Lupinus perennis</i> (H) **	Sundial Lupine	Early Summer	Blue-Violet	Moist to Dry well drained sandy	14"-30"/12"
<i>Marshallia grandiflora</i>	Barbara's Buttons	Early Summer	Rose Pink to Lavendar	Wet to Moist	10"-14"/12"
<i>Medeola virginiana</i> (H)***	Indian Cucumber-root	Late Spring - Early Summer	Yellowish Green	Moist	1'-2'/1'
<i>Mimulus alatus</i> (H)	Winged Monkey-flower	Summer	Pinkish, Purpleish	Wet to Moist	1'-3'/1'
<i>Mimulus ringens</i> (H)	Allegheny Monkey-flower	Summer	Blueish Violet	Wet to Moist	2'-4'/1'
<i>Monarda clinopodia</i> (H)	Basil Bee Balm	Summer	White	Moist	2'-4'/18"
<i>Monarda didyma</i> (H)	Bee Balm, Oswego Tea	Mid- Summer	Red	Wet to Moist	3'-4'/2'-3'
<i>Monarda fistulosa</i> (H)	Wild Bergamot	Mid-Summer	Lavender	Moist to Dry	3'-4'/2'-3'
<i>Monarda media</i> (H)	Purple Bergamot	Summer	Fushia to	Moist	2'-3'/18"

## Native Herbaceous Plants for Sun to Part Sun

Manada Native Plant Database					
Scientific Name	Common name	Bloom Time/Fruiting Time	Bloom Color	Soil Moisture	Height
			Purple		
<i>Oenothera fruticosa</i> (H)	Sundrops	Late Spring	Yellow	Moist to Dry	1.5'/1.5'
<i>Penstemon digitalis</i> (H)	Tall White Beardtongue	Spring	White	Moist	2'-4'/1.5'
<i>Penstemon hirsutus</i> (H) ***	Northeastern Beardtongue, Hairy Beardtongue	Early Summer	Violet to Pale Rose	Moist to Dry, well-drained	12"-18"/12"
<i>Penstemon laevigatus</i> (H)	Eastern Beardtongue	Late Spring - Mid Summer	Light Purple to White	Moist to Dry	1'-4'/1'
<i>Phlox maculata</i> (H)	Meadow Phlox	Summer (before Garden Phlox)	Pinkish-purple	Wet to Moist	2'-3'/2'
<i>Phlox paniculata</i> (H)	Garden Phlox, Summer Phlox	Summer	Pinkish-purple or white	Moist to Dry	3'-5'/2'-3'
<i>Phlox pilosa</i> (H)	Downy Phlox, Prairie Phlox	Late Spring	Pinkish-purple	Moist to Dry	1'-2'/1'-1.5'
<i>Phlox sublata</i> (H)	Moss-pink, Moss Phlox	Spring	Lavender, Pink, White	Well-drained	2"-6"/14"-18"
<i>Physostegia virginica</i> (H)	Obedient plant	Late Summer	Light to Dark Pink, White	Wet to Moderately Dry	2'-5'/3-4'
<i>Pycnanthemum</i> (various species) (H)	Mountain Mints	Summer into fall	White	Moist to Average	3'-4'/2'-3'
<i>Ratibida pinnata</i> (H) **	Prairie Coneflower, Grey-headed Coneflower, Grey-headed Coneflower	Summer	Yellow with brown center	Moist to Dry well drained	2'-4'/2'
<i>Rhexia virginica</i> (H)	Virginia Meadow Beauty	Mid Summer to Late Summer	Rose Pink	Wet to Moist	8"-28"/12"-16"
<i>Rudbeckia fulgida</i> (H)	Black-eyed Susan	Summer	Yellow	Moist to Dry	2'-2.5'/3'
<i>Rudbeckia fulgida</i> var. <i>fulgida</i> (H)	Showy Coneflower, Eastern Coneflower, Black-eyed Susan	Late Summer-Fall	Yellow	Moist to Dry	2' - 2.5'
<i>Rudbeckia hirta</i> (H)	Black-eyed Susan	Summer	Yellow with black Center	Moist to Dry	2'-3'/1'-1.5'
<i>Rudbeckia lacinata</i> (H)	Cutleaf Coneflower, Green-headed Coneflower	Summer	Yellow	Wet to Moist to Dry	3'-6'/3'-4'

## Native Herbaceous Plants for Sun to Part Sun

Manada Native Plant Database					
Scientific Name	Common name	Bloom Time/Fruiting Time	Bloom Color	Soil Moisture	Height
Rudbeckia triloba (H)	Brown-eyed Susan, Three-lobed Coneflower	Summer	Yellow	Moist to Dry	3'-4'/2'
Ruellia caroliniensis (H)	Carolina Petunia, Hairy Petunia	Summer	Lavender-blue	Moist to Dry	1'-2.5'/18"
Ruellia humilis (H)	Fringeleaf Petunia	Summer	Light Blue	Moist to Dry	2'-3'/1'
Ruellia strepens (H)	Limestone Petunia	Spring into Summer	Lavender-blue	Moist to Dry	1'-3'/18"
Salvia lyrata (H)	Lyre-leafed Sage	Spring - Early Summer	Light Purple	Wet to Dry	1'-2'/1'
Scutellaria incana (H)	Downy Skullcap	Mid - Late Summer	Violet Blue	Moist to Average	2'-3'/2'
Scutellaria integrifolia (H) **	Hyssop Skullcap	Mid-summer	Violet Blue	Wet, Moist to Dry, well-drained	16"-20"/8"-12"
Scutellaria serrata (H)	Allegheny Skullcap	Early Summer	Violet Blue	Moist to Dry	14"-20"/18"
Sedum telephiodides (H)	Allegheny Stonecrop, Wild Live-forever	Late Summer to Fall	Pale Pink	Moist, well-drained	8"-16"/8"-12"
Sedum ternatum (H)	Wild Stonecrop	Spring	White	Moist to Average	4"-8"/8"-12"
Senecio aureas (H)	Golden Ragwort	Spring	Yellow	Moist, good drainage	1'/1'
Senna hebecarpa (H)	Wild Senna	Summer	Yellow	Moist	'-6'
Silene caroliniana (H)	Sticky Catchfly, Wild Pink	Late Spring	Light to Dark Pink	Moist to Dry	4"-8"/8"
Silene virginica (H)	Fire Pink, Catchfly	Summer	Scarlet to Crimson	Moist, Well-drained	8"-14"/12"
Silphium trifoliatum var. trifoliatum (H)	Wholed Rosinweed	Mid-Summer	Yellow	Moist to Dry	3'-4'/2'-3'
Sisyrinchium angustifolium (H)	Blue-eyed Grass	Late Spring to Early Summer	Light Blue	Moist	12"-20"/6"-8"
Solidago caesia (H)	Blue-stemmed Goldenrod	Fall	Yellow	Moist, Good Drainage	1.5'-4'/1.5'
Solidago juncea (H) **	Early Goldenrod	Summer	Yellow	Dry	1.5'04'/2'
Solidago nemoralis (H)	Gray Goldenrod	Fall	Yellow	Dry, well-drained	1'-2'/1'
Solidago odora (H)	Sweet Goldenrod	Late Summer	Yellow	Moist to Dry	2'-5'/2'
Solidago rigida (H) **	Stiff Goldenrod	Late Summr to Fall	Yellow	Moist	3'-5'/1'-2'
Solidago rugosa	Wrinkle-leaf goldenrod,	Fall	Yellow	Moist to	2'-6'/3'

## Native Herbaceous Plants for Sun to Part Sun

Manada Native Plant Database					
Scientific Name	Common name	Bloom Time/Fruiting Time	Bloom Color	Soil Moisture	Height
"Fireworks" (H)	Rough-stemmed Goldenrod			Average	
<i>Solidago speciosa</i> (H)	Showy Goldenrod	Late Summer - Fall	Yellow	Moist to Dry	2'-5'/3'
<i>Thalictrum pubescens</i> (H)	Tall Meadow Rue	Summer	White	Wet to Moist	3'-10'/2'
<i>Tradescantia ohiensis</i> (H)	Spiderwort	Summer	Blue	Moist to Dry, well-drained	24"-30"/12"
<i>Tradescantia virginiana</i> (H)	Virginia Spiderwort	Spring	Blue, Violet	Moist	1'-2'/2'
<i>Trichostema dichotomum</i> (H)	Forked Blue-curls	Summer into Fall	Deep Blue	Moist to Dry	4"-30"
<i>Trollius laxus</i> (H)	Spreading Globeflower	Early Spring into Early summer	Yellow	Wet to Moist	12"-20"/12"-16"
<i>Verbena hastata</i> (H) **	Blue Vervain	Summer	Blue-Violet	Wet to Moist	2'-4'/1'-1.5'
<i>Vernonia glauca</i> (H) **	Upland ironweed	Late Summer	Purple	Moist to Dry	3'-5'/2'-4'
<i>Vernonia novaboracensis</i> (H) **	New York ironweed	Late Summer	Purple	Wet to Moist	4'-7'/2'-4'
<i>Veronicastrum virginicum</i>	Culver's root	Summer	White	Wet, Moist to Average	4'-5'/2'-3'
<i>Zizia aptera</i> (H)	Heart-leaved Alexander	Late Spring	Yellow	Moist to Average	1'-2'/1.5'
<i>Zizia aurea</i> (H)	Golden Alexander	Late Spring	Yellow	Wet to Moist	1'-3'/1.5'

\*\* Full Sun

\*\*\* Part Sun